

Gujarat Groundnut Acreage, Yield and Production Estimates (Summer-2023)

| Sr. No. | District | Groundnut Acreage 2022 (ha) | Groundnut Acreage 2023 (ha) | Yield 2022 (kg/ha) | Yield 2023 (kg/ha) | Production 2022 (Tonnes) | Production 2023 (Tonnes) |
|--------------------|--------------|-----------------------------|-----------------------------|--------------------|--------------------|--------------------------|--------------------------|
| 1 | Banaskantha* | 24,000 | 25,195 | 1812 | 1943 | 43,488 | 48,954 |
| 2 | Bhavnagar | 8,000 | 8,300 | 1889 | 1940 | 15,112 | 16,102 |
| 3 | Amreli | 3,500 | 2,900 | 1697 | 1529 | 5,940 | 4,434 |
| 4 | Rajkot | 1,900 | 1,800 | NA | 1480 | NA | 2,664 |
| 5 | Sabarkantha | 1,900 | 1,500 | NA | 1578 | NA | 2,367 |
| Sub Total | | 39,300 | 39,695 | | | 64,540 | 74,521 |
| Others | | 20,700 | 14,200 | | | 41,090 | 26,828 |
| State Total | | 60,000 | 53,895 | 1,796 | 1,880 | 1,05,629 | 1,01,349 |

- Source of the district level acreage data is DES (<https://eands.dacnet.nic.in/>).
- * Remote sensing based estimated acreage.
- Last year Junagarh and Jamnagar were surveyed instead of Rajkot and Sabarkantha as area has reduced in these districts this year.
- Production has been calculated as *Acreage x Questionnaire Based Yield Data*. Highest Summer Groundnut Yield has been recorded in Banaskantha district and lowest in Rajkot District.
- Summer Groundnut sown acreage has decreased by 10% as compare to last year due to increased Bajra acreage in the state.
- On the other hand production has reduced by 4% only due to higher yield recorded in Banaskantha and Bhavnagar district. However the yield is low in Amreli, and Rajkot which can be attributed to unseasonal rainfall in the month of April and May, 2023.